



City Research Online

City, University of London Institutional Repository

Citation: McManus, S., Bebbington, P. E., Tanczer, L., Scott, S. & Howard, L. M. (2023). Receiving threatening or obscene messages from a partner and mental health, self-harm, and suicidality: results from the Adult Psychiatric Morbidity Survey. *Social Psychiatry and Psychiatric Epidemiology: the international journal for research in social and genetic epidemiology and mental health services*, 58(12), pp. 1749-1759. doi: 10.1007/s00127-021-02113-w

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/26529/>

Link to published version: <https://doi.org/10.1007/s00127-021-02113-w>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk



Receiving threatening or obscene messages from a partner and mental health, self-harm and suicidality: results from the Adult Psychiatric Morbidity Survey

Sally McManus^{1,2} · Paul E. Bebbington³ · Leonie Tanczer⁴ · Sara Scott⁵ · Louise M. Howard⁶

Received: 28 December 2020 / Accepted: 2 June 2021
© The Author(s) 2021

Abstract

Purpose Threatening or obscene messaging is repeated, unwanted texts, emails, letters or cards experienced by the recipient as threatening or obscene, and causing fear, alarm or distress. It is rarely examined as an aspect of intimate partner violence. We describe the prevalence of exposure to threatening/obscene messaging from a current or ex-partner; characteristics of victims; and associations with other forms of violence and abuse, mental disorder, self-harm, and suicidality.

Methods Cross-sectional probability-sample survey of the general population in England aged 16+ . Multivariable regression modelling tested associations between receipt of threatening/obscene messaging and current common mental disorder, past-year self-harm and suicidality.

Results Threatening/obscene messages were received from a current/ex-partner by 6.6% (95%CI: 5.9–7.3) of adults who had been in a relationship; 1.7% received these in the past year. Victims were more likely to be female, under 35, single or divorced, socioeconomically disadvantaged, and to have experienced other forms of sexual and partner violence and abuse. Those who received threatening/obscene messages in the past year were more likely to experience common mental disorder (adjusted odds ratio 1.89; 1.01–3.55), self-harm (2.31; 1.00–5.33), and suicidal thoughts (2.00; 1.06–3.78).

Conclusion Threatening/obscene messaging commonly occurs in the context of intimate partner violence. While often occurring alongside sexual and physical violence, messaging has an additional association with mental disorder and suicidality. Routine enquiry in service settings concerning safety, including those working with people who have escaped domestic violence, should ask about ongoing contact from previous as well as current partners. This should include asking about messaging, as well as other forms of potentially technology-enabled abuse which may become increasingly common.

Keywords Intimate partner violence · Emotional abuse · Cyber bullying · Threat · Mental health · Self-harm · Suicidality

This article belongs to the Topical Collection Violence and Women's Mental Health.

✉ Sally McManus
sally.mcmanus@city.ac.uk

¹ Violence and Society Centre, City, University of London, London, UK

² National Centre for Social Research, 35 Northampton Square, London EC1V 0AX, UK

³ Psychiatry Division, University College London, London, UK

⁴ Faculty of Engineering Science, University College London, London, UK

⁵ DMSS, North Dalton, England, UK

⁶ Section of Women's Mental Health, Institute of Psychology, Psychiatry and Neuroscience, Kings College London, London, UK

Introduction

Violence, abuse, and bullying are widely recognised as factors contributing to both the onset and continuation of mental disorder [1, 2]. England's Department of Health and Social Care (DHSC) recommends that healthcare professionals ask patients about current and historic exposure to violence and abuse, to identify risk factors, establish safety, and support recovery [3–5]. A dose–response relationship has emerged with the risk of poor mental health being greater where experience of violence and abuse has been more pervasive [6]. People exposed to one type of violence and abuse are more likely to also experience other types, at the same time or subsequently [7].

One of the most common forms of violence in adulthood is that from an intimate partner (IPV). The World Health

Organisation lists different types of IPV, including physical violence, sexual violence, emotional or psychological abuse, and controlling behaviours, involving isolation, monitoring, and restricted access to resources [8]. In the UK, the *Serious Crime Act 2015* was the first to recognise controlling or coercive behaviour in the context of an intimate or family relationship as an offence [9], and the *Domestic Abuse Act 2021* specifically recognises economic and technology-related abuse [10, 11]. However, a tension has been noted between such public policy and the gendered reality of domestic violent crime [12].

Emotional abuse is known to predict poor mental health as strongly as physical or sexual abuse [13] and includes insults, belittling, intimidation and threats, delivered face to face or communicated in other ways. The *Malicious Communications Act 1988* had made it illegal in England and Wales to send or deliver letters or other articles for the purpose of causing distress or anxiety, and in 2001 the legislation was updated to also apply to electronic communications [14]. From March 2019 to March 2020, 36% of all domestic violence-related stalking and harassment offences recorded by police in England and Wales were malicious communications [15]. Police forces, however, have been found to often not record—or to mis-record—such reports [16]. Crime statistics therefore significantly underestimate the extent of abusive behaviours conducted through communications, including digital communications, in the population, with further changes to likelihood of reporting occurring in the context of COVID-19 restrictions [17].

Global developments in communication technology have the potential to enable malicious communications to become increasingly pervasive, immediate, and intrusive [18]. While the UK government's recent *Online Harms* white paper acknowledged this, it made no reference to its occurrence in the context of IPV nor accounted for the growing risk from Internet-connected devices [19]. Instead attention has focussed on threatening or obscene messaging in other situations: such as hate speech [20], children's online behaviour [21], and conduct on social media. Furthermore, research on sexting [22] and cyberbullying has tended to focus on peer-related bullying rather than IPV. Malicious communications can form an aspect of technology-facilitated abuse ('tech abuse'), which includes the monitoring of victims through networked camera systems, stalking via software and tracking tools frequently referred to as "stalkerware" [23, 24], and other forms of "smart" technological systems such as Internet-connected household appliances to coerce, control, and harm others [25]. Limited recognition has been given to these newer forms of abuse, especially in the context of emotional abuse and coercive control from a current or previous partner. In 2020, the UK Law Commission [26] launched a consultation on reforming the Malicious Communications Act 1988 and Communications Act 2003, in light of

developments in online communication and recognising the inherently gendered nature of online abuse and its potential for causing harm.

While England's mental health survey did not ask specifically about tech abuse, it did include questions on receipt of repeated threatening and obscene messages from a current or previous partner, spanning both technology-facilitated and other forms of communication. The aims of this study were to establish:

- The first prevalence estimates of exposure to threatening/obscene messaging from current or previous partners among adults in England,
- The characteristics of people exposed to such messages at any point in adulthood, and specifically within the past year,
- Whether past-year exposure to such messaging was associated with common mental disorder, non-suicidal self-harm, suicidal thoughts, and suicide attempt, after adjustment for demographic and socioeconomic circumstances and experience of other forms of violence and abuse.

Methods

We carried out a secondary analysis of data from the 2014 Adult Psychiatric Morbidity Survey (APMS), a general population survey of the mental health of adults resident in private households in England. A stratified random probability sampling design was used, based on the selection of addresses from the Postcode Address File (PAF). This involved multiple stages: sampling primary sampling units (PSUs); addresses within selected PSUs; and one individual aged 16 or over from each selected address. People living in communal or institutional establishments, in temporary housing, or sleeping rough, were not in scope. Fieldwork took place May 2014–September 2015, with verbal informed consent.

Seven thousand five hundred forty-six participants were interviewed, a response rate of 57%. Weights were developed to take account of selection probabilities and non-response, in order to render results representative of the household population. Interviews averaged 1.5 h and were conducted in people's own homes (or elsewhere, if preferred) by trained research interviewers. The questionnaire was largely administered face-to-face using computer-assisted interviewing. Some information considered more sensitive was collected by self-completion for increased privacy, with participants keying their responses directly into a laptop. Interviewers assisted respondents with literacy or eyesight difficulties. 7058 participants carried out the self-completion component: factors associated with declining the self-completion included being male and being older [27].

Further methodological details are published elsewhere [28, 29].

Measures

Exposure

The receipt of repeated threatening or obscene messages was established by asking: ‘Has a partner or ex-partner ever sent you more than one unwanted letter, email, text message or card that was either obscene or threatening and which caused you fear, alarm or distress?’ On- and offline messaging could not be disaggregated, nor whether messages were experienced as either threatening or obscene. The question was administered by self-completion as part of a module about violence and abuse from a partner or ex-partner. It was only asked of those who had had a partner (boyfriend/girlfriend/spouse) at some point. Follow-up questions established how recently the abuse occurred. If the experience had happened in the past 12 months, respondents were asked about frequency.

Outcomes

Common mental disorders (CMDs) were assessed using the Clinical Interview Schedule-Revised (CIS-R). This is an interviewer-administered structured interview covering the presence of non-psychotic symptoms in the week prior to interview. It can provide prevalence estimates for six CMDs according to ICD-10 clinical criteria [30]: generalised anxiety disorder, phobia, obsessive compulsive disorder, panic disorder, depression and other common mental disorder not otherwise specified [31]. The outcome variable was binary, indicating either presence or absence of any CMD.

Suicidal thoughts: In the face-to-face section of the interview, participants were asked: ‘Have you ever thought of taking your life, even though you would not actually do it?’ An affirmative response was followed with a question about when this had last occurred, and a variable was derived indicating those reporting such thoughts in the past year.

Non-suicidal self-injury and suicide attempt: the 5th Diagnostic and Statistical Manual of Mental Disorders includes non-suicidal self-injury and suicidal behaviour disorder as conditions for further study [32]. Suicide attempts and non-suicidal self-harm were examined separately. Questions about suicide attempts within the past year were asked in both the face-to-face and self-completion sections of the interview: ‘Have you ever made an attempt to take your life, by taking an overdose of tablets or in some other way?’ A derived variable combined reports of a suicide attempt in the past year in either section of the interview. Non-suicidal self-harm was also asked both face-to-face and

in the self-completion section: ‘Have you ever deliberately harmed yourself in any way but not with the intention of killing yourself?’ Non-suicidal self-harm in the past year similarly drew on reports from either the face-to-face or self-completion section. While agreement was high, rates from the self-completion section were higher [29].

Harmful or dependent alcohol use: was measured using the Alcohol Use Disorders Identification Test (AUDIT) [33]. The AUDIT takes the year before the interview as a reference period, consists of 10 items and covers alcohol consumption, alcohol-related harm, symptoms of alcohol dependence. Answers to all questions were scored from zero to four, and summed to give a total score ranging from 0 to 40. A binary indicator was produced for scores of 1–15 and 16 or over, with the latter indicative of a level of alcohol use considered harmful drinking and/or dependent.

Covariates

Emotional, physical and sexual violence and abuse in childhood and adulthood were asked about in the self-completion section of the interview. Those relating to violence by a current or former partner were adapted from questions in the British Crime Survey, originally based on the Conflict Tactics Scale (CTS) [34]. Emotional abuse in childhood was indicated by an affirmative response to: ‘Before you were 18, did you get scared or feel really bad because an adult in your life called you names, said mean things to you, or said they didn’t want you?’ Physical violence in childhood was based on: ‘Not including smacking, before you were 18, did an adult in your life hit, beat, kick, or physically hurt you in any way?’ Childhood sexual abuse was derived from two questions relating to non-consensual sexual contact and forced sexual intercourse before age 16. Emotional abuse from a current or former partner was based on an affirmative response to: ‘Has a partner or ex-partner ever repeatedly belittled you to the extent that you felt worthless?’ Physical violence from a partner was established by asking: ‘Has a partner or ex-partner ever pushed you, held or pinned you down or slapped you?’ and ‘Has a partner or ex-partner ever kicked you, bit you, or hit you with a fist or something else, or threw something at you that hurt you?’ Sexual violence or abuse in adulthood was derived from two questions about non-consensual sexual contact and forced sexual intercourse since age 16.

Standard demographic questions established sex (male, female), age (banded into 16–24, 25–34, 35–44, 45–54, 55+) and marital status (single; married/cohabiting; separated/divorced/widowed). Ethnicity was self-ascribed, and, due to the relatively small number of participants in the White Other, Asian/Asian British, Black, Mixed and Other groups, grouped into White British and Other. Socio-economic context was captured from participants’ housing

tenure (owner-occupier, renting from a social landlord, renting from a private landlord) and employment status (employed, unemployed and looking for work, economically inactive). Household income was equivalised (to take account of number and ages of people living in the household), and analysed in quintiles. The questionnaire and further methodological details are published elsewhere [29].

Statistical analysis

Analyses used weighted data and took account of the complex survey design (sample stratification and clustering) and non-response. Data management, descriptive and correlational analyses were conducted in SPSS v25 [35], with *p*-values and confidence intervals calculated at the 95% level. Stata v14.1 [36] was used for running multiple variable logistic regression analyses. Regression models were run for four binary-coded dependent variables: current CMD; past-year suicidal thoughts; past-year suicide attempts; and past-year non-suicidal self-harm. Each was analysed according to three sequential models. The first examined past-year exposure to threatening/obscene messages as an independent variable, controlling for potential demographic and socioeconomic confounders. The second model additionally controlled for exposure to forms of violence and abuse other than emotional abuse in adulthood; the final model additionally controlled for emotional abuse in adulthood as well. Multicollinearity was checked, and found not to be a problem, with the Variance Inflation Factor (VIF) for all variables below 2. Missing data were minimal for all variables and excluded from analyses, except for household income. Household income was not provided by 23.4% of the sample. It is likely that this was either because participants did not know the income of all household members or because income is considered sensitive to report. For this variable, therefore, those not providing a response were coded as 'missing' and retained in the analysis.

Results

Our results suggest that in England around 96% ($n=6857$) of adults aged 16 or over have had at least one partner or spouse. Men (5.4%, $n=122$) were more likely than women (2.7%, $n=79$) to have never had a partner ($p<0.001$). Subsequent analyses are based on participants reporting at least one partner in their life, to avoid results being confounded by relationship history.

Prevalence

One person in fifteen (6.6%; $n=484$; 95% CI 5.9–7.3) had received two or more unwanted obscene or threatening texts, emails, letters, or cards, that had caused them fear, alarm or distress from a current or former partner. This was twice as likely in women (8.7%) as men (4.4%) ($p<0.001$). 14.5% of women aged 16–24 reported receipt of threatening/obscene messages, three times the rate for men of the same age (4.3%). A quarter (25.1%) of people exposed to threatening/obscene messages had received more than one in the past year (1.7% of adults, $n=124$). Two-fifths (40.1%) of those with past-year exposure had received such messages monthly or more often.

Characteristics

Table 1 shows that people who had been exposed to threatening/obscene messages from a partner were more likely: to be young (aged 16–24 (18.4%) or 25–34 (28.7%), $p=0.005$), single or divorced (52.7%, $p<0.001$), unemployed (5.3%, $p<0.001$), and living in rented accommodation (58.0%, $p<0.001$) and lower income households (18.7%, $p=0.002$). While associated with disadvantage, the experience of threatening/obscene messaging was evident in all groups. The profile of those exposed in the past year was similar to that for those who had ever been exposed, although recent experience was more likely in young. There was no significant association with ethnic group ($p=0.505$), although the sample was too small to examine this definitively.

People who received threatening/obscene messages from a current or former partner were more likely to have also experienced all the other types of violence and abuse examined. They were three times more likely than those who had not experienced such messaging to have been abused in childhood emotionally (30.2%, cf. 8.9%), sexually (23.5%, cf. 6.8%), and/or physically (26.5%, cf. 11.4%). They were also about five times more likely to have experienced emotional (59.4%, cf. 8.8%) or physical abuse from a partner (63.1%, cf. 11.4%), or sexual abuse (22.0%, cf. 4.4%) at some point in adulthood. Two-thirds (69.7%) of women and half of men (48.8%) who received threatening/obscene messages had experienced physical partner-violence at some point in adulthood, compared with 14.6% of women and 8.2% of men who had not received threatening/obscene messages.

Rates of CMD were more than twice as high in people who had received threatening/obscene messages (39.2%) than in those who had not (15.2%). This pattern was evident both in women (42.8%, cf. 18.3%) and men (31.5%, cf. 12.0%). Non-suicidal self-harm rates were higher in women (7.1%) and men (3.0%) exposed to such messages, than in women (1.7%) and men (1.3%) not exposed. Suicidal

Table 1 Demographic and socioeconomic profile of those exposed and not exposed to repeated threatening/obscene messaging from a partner or ex-partner

Characteristics	Total ^a		Past year exposure		Ever exposed (including past year)		Never exposed		p-value ^b
	N	Weighted %	n	Weighted %	n	Weighted %	n		
Total	6804	100	124	1.7	484	6.6	6320	93.4	
Sex									
Men	2731	48.3	41	45.8	114	32.1	2617	49.4	<0.001
Women	4073	51.7	83	54.2	369	67.9	3704	50.6	
Age									
16–24	461	12.5	18	24.3	53	18.4	408	12.1	0.005
25–34	971	17.5	43	33.0	129	28.7	842	16.7	
35–44	1099	16.6	27	17.8	106	20.0	993	16.4	
45–54	1213	18.2	25	19.5	110	20.0	1103	18.1	
55 +	3060	35.2	11	5.4	86	12.9	2974	36.8	
Ethnic group									
White British	5824	82.3	101	78.5	413	83.6	5411	82.2	0.505
Other	963	17.7	23	21.5	69	16.4	894	17.8	
Marital/cohabitation status									
Married/cohabiting	3896	64.6	25	29.7	181	47.3	3715	65.9	<0.001
Single	1301	21.7	61	50.5	165	34.7	1136	20.7	
Divorced	1607	13.7	38	19.8	138	18.0	1469	13.4	
Economic activity									
Employed	3745	61.5	82	66.5	316	69.9	3429	60.9	<0.001
Unemployed	189	3.1	7	6.1	23	5.3	166	2.9	
Other	2870	35.4	35	27.3	145	24.8	2725	36.2	
Tenure									
Owner occupied	4510	65.2	42	65.6	192	42.0	4318	66.8	<0.001
Social renter	1084	15.0	30	14.8	134	26.0	950	14.2	
Private or other	1177	19.9	51	19.6	154	32.0	1023	19.0	
Equalised household income quintiles									
Highest income	1072	16.4	11	7.5	50	11.0	1022	16.7	0.002
2nd	1037	15.2	19	13.7	73	16.6	964	15.2	
3rd	1151	15.8	28	21.3	88	16.7	1063	15.8	
4th	1145	15.0	23	20.9	84	15.6	1061	15.0	
Lowest income	1057	14.1	32	20.9	113	18.7	944	13.8	
Missing	1342	23.4	10	15.8	76	21.3	1266	23.6	

^aTotal: adults aged 16 or over reporting having had at least one partner (girlfriend/boyfriend/spouse)^bAssociation between characteristic and threatening or obscene messaging ever

thoughts were three times more common in those exposed to messages (12.6%) than in those who were not (4.1%), while attempted suicide was four times more likely (2.0%, cf. 0.5%). Exposure to abusive messaging was also associated with harmful or dependent use of alcohol. For each of these outcomes, the pattern of association was similar for women and men (see Table 2).

Adjusted associations

In a model controlling for demographic and socioeconomic factors, people exposed to threatening/obscene messages in the past year had 3.95 times the odds of CMD compared with the rest of the population (95% CI 2.5–6.3, $p < 0.001$). After further adjustment for exposure to physical violence (in childhood and adulthood), sexual abuse (in childhood and adulthood), and emotional abuse (in childhood only), the association between recent threatening/obscene messages and CMD was attenuated but remained significant (adjusted OR [aOR] 2.20, 1.2–4.0, $p = 0.011$). Past-year messaging retained an independent association

with CMD even when further controlling for experience of any form of emotional abuse from a partner at any point in adulthood (aOR 1.89, 1.0–3.5, $p = 0.047$), see Table 3 and supplementary materials.

Controlling for demographic and socioeconomic factors, those with past-year exposure to threatening/obscene messages had odds of non-suicidal self-harm in the past year 4.28 times higher than the rest of the population (2.0–9.3, $p < 0.001$). Further adjustment for physical violence and sexual abuse as a child and adult, as well as for childhood emotional abuse, reduced the odds ratio to 2.95 (1.2–7.1, $p = 0.015$). In the final model, which additionally adjusted for emotional abuse from a partner across adulthood, the effect size was attenuated (aOR 2.31), although the independent association between past-year threatening/obscene messages and non-suicidal self-harm was not significant at the 95% level (1.0–5.3, $p = 0.051$).

The odds of having suicidal thoughts in the past year remained 3.76 times higher in those exposed to messages than in those not exposed, when adjusting for demographic and socioeconomic factors (2.1–6.6, $p < 0.001$). In a model

Table 2 Prevalence of other types of violence and abuse, mental disorder, self-harm, suicidality and harmful/dependent alcohol use by whether exposed to threatening/obscene messages from a (ex)partner

	Men			Women			Total		
	Threatening/ obscene mes- sages	Not exposed	Total	Threatening/ obscene mes- sages	Not exposed	Total	Threatening/ obscene mes- sages	Not exposed	Total
Type of violence and abuse ^a	%	%	%	%	%	%	%	%	%
Physical abuse from caregiver in childhood	31.9	13.4	14.2	24.0	9.5	10.8	26.5	11.4	12.4
Sexual abuse in childhood	14.4	4.0	4.5	27.7	9.6	11.1	23.5	6.8	7.9
Emotional abuse from caregiver in childhood	33.8	7.7	8.8	28.6	10.1	11.7	30.2	8.9	10.3
Physical violence from partner in adulthood	48.8	8.2	10.0	69.7	14.6	19.3	63.1	11.4	14.8
Sexual abuse or violence in adulthood	6.8	1.5	1.7	29.1	7.3	9.2	22.0	4.4	5.6
Emotional abuse from partner in adulthood	42.1	4.9	6.6	67.6	12.7	17.4	59.4	8.8	12.2
Mental disorder and self-harm									
Common mental disorder (CMD)	31.5	12.0	12.9	42.8	18.3	20.5	39.2	15.2	16.8
Non-suicidal self-harm in past year	3.0	1.2	1.3	7.1	1.7	2.2	5.8	1.5	1.7
Suicidal thoughts in past year	13.0	4.4	4.8	12.4	3.8	4.6	12.6	4.1	4.7
Suicide attempt in past year	1.3	0.5	0.5	2.4	0.5	0.7	2.0	0.5	0.6
Harmful/dependent alcohol use	12.0	4.4	4.7	5.8	1.5	1.8	7.8	2.9	3.2
N	114	2617	2731	370	3703	4073	484	6320	6804

^aCMD, suicidal thoughts, suicide attempt, non-suicidal self-harm and experience of all types of violence and abuse were more likely in those who had received threatening/obscene messages in the past year at $p < 0.001$

	Models 1 ^a				Models 2 ^b				Models 3 ^c			
	Adj. OR	Lower CI	Upper CI	<i>p</i> -value	Adj. OR	Lower CI	Upper CI	<i>p</i> -value	Adj. OR	Lower CI	Upper CI	<i>p</i> -value
Common mental disorder	3.95	2.48	6.31	<0.001	2.20	1.20	4.05	0.011	1.89	1.01	3.55	0.047
Non-suicidal self-harm in past year	4.28	1.98	9.27	<0.001	2.95	1.23	7.07	0.015	2.31	1.00	5.33	0.051
Suicidal thoughts in past year	3.76	2.13	6.63	0.000	2.19	1.13	4.24	0.021	2.00	1.06	3.78	0.034
Suicide attempt in past year	5.49	2.02	14.98	0.001	2.94	0.86	10.07	0.086	2.35	0.77	7.16	0.132
Harmful/dependent alcohol use	2.05	0.89	4.73	0.090	1.15	0.51	2.60	0.744	1.00	0.45	2.24	0.996

Models 2: model 1 plus five indicators of violence and abuse; sexual and physical abuse in childhood or adulthood, and emotional abuse in childhood

Models 3: model 2 plus emotional abuse in adulthood

People who received repeated threatening or obscene messages from a current or previous partner in the past year were more likely to experience anxiety and depression, to engage in self-harm, and to be suicidal. These associations remained elevated even after adjusting for confounding for current socioeconomic disadvantage and exposure to a

range of types of violence and abuse across the life course. Lindsay and colleagues [42] have shown how abusive messaging from an intimate partner increases the odds of experiencing fear as a result of victimisation, and that, in turn, fear was associated with an increased risk of depression. Our results are consistent with a trauma-informed understanding of mental health, and support the evidence base by demonstrating the additional effect that such communications can have. Health care professionals carrying out safety assessments should therefore enquire routinely and explicitly about messages and communications, and these should refer to any received from a previous as well as a current partner. Voluntary and statutory support services engaging with people who have left abusive relationships should establish whether survivors still have ongoing contact with former partners, and the nature of any communication, to ensure threats, intimidation, harassment and monitoring are not missed.

This study was unable to examine technology-enabled abuse in isolation. To respond to emerging risks—with implications for the frequency, immediacy, and visual content of threatening or obscene messaging—technology-enabled communications should be asked about in research and during safety assessments and accounted for during safety planning. New telephone numbers, contact details, and ideally devices, should be issued to those at risk [43]. Mobile providers and other suppliers should facilitate blocking other people across platforms. As technology may change the ways in which abuse is mediated, future surveys will need to adapt to ensure that the scale and impact of such experiences in adulthood and childhood is captured, including on longitudinal studies. The rapidity of such changes poses challenges for research, for service responses, and for victims.

This is the first analysis using representative English mental health survey data to present the national prevalence of threatening and obscene messaging from partners, together with its social and mental health correlates. The survey was based on a national probability sample and high-quality self-completion procedures in the context of a face to face interview. The identification of threatening or obscene messaging, however, was based on a single question, which did not distinguish between online and offline forms of communication, whether the perpetrator was a current or former partner, or whether messages were felt to be threatening or obscene. In the context of what Herman [44] calls ‘domestic terrorism’, however, sexual comments and pornographic images from a current or previous partner will often be experienced by women as *both* threatening and obscene. The question also asked only about the receipt of threatening or obscene messages that had actually provoked fear, distress, or alarm in the participant: in consequence, it failed to capture people who had received such messages but felt unaffected by them. The lower rate in men, for example,

might have reflected the fact they were less likely to experience distress from such messages, or were less likely to disclose having felt distress. Recent (past year) experience was reported by relatively few (124) participants, and some of the analyses were accordingly underpowered. The findings, however, echo gendered dynamics identified in the context of the sending and receiving of sexualised images. Female teens receive far more unsolicited messages and images than males and are also more stigmatised for sending, for example, nude pictures in return [45]. The Conflict Tactics Scale (CTS) [34], which informed the questions used to capture other forms of partner violence and abuse, has been identified as problematic. The scale amalgamates non-equivalent types of gendered violence (for example, implying that being ‘slapped’ and ‘pinned down’ are equivalent) [46], and fails to capture context or impact [47].

The sampling frame excluded those living in residential or institutional settings or sleeping rough; and people living in domestic violence refuges and shelters were unlikely to have been selected. However, the proportion of the population not living in private households is small [48] and would have had little impact on the prevalence estimates and associations examined. While a survey response rate of 57% is in line with similar surveys, bias due to non-participation is inevitable. While cross sectional surveys using high quality general population samples are the best data source for prevalence and for profiling (understanding nuanced socioeconomic circumstances and the relationship with other types of violence and abuse), conclusions cannot be drawn about causality.

Conclusion

This study supports earlier research in finding pronounced associations between emotional abuse and poor mental health and suicidality. When asking about violence and abuse, clinicians, law enforcement, and service providers need to ensure specific enquiry about threats, intimidation, and harassment from partners who may not be present in the home. Such enquiry, as well as future research, must address the potential for ongoing abuse from former partners, who may use remote communications to continue to exert threats and control [37]. Abusive messaging is one strategy of control that partners, especially former partners, use when their victim is not physically available [49]. In the context of the unintended impacts of ongoing physical distancing measures to mitigate against Covid-19 transmission [50–52], addressing abusive messaging has only gained in urgency.

Mental health services and future research should cover both online and offline modes of communication. Malicious communications in the context of IPV should feature in policy and legislation on both domestic violence

and online harms. Our results support calls for technology companies to publish annual transparency reports to assess the prevalence of harmful content on their platforms. Hopefully through such means more data and evidence on online and offline facilitated forms of communication-enabled abuse will become available and will help to contextualise future research in this area.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00127-021-02113-w>.

Author contributions SM designed and conducted the analyses and drafted the manuscript. PEB, SS, LT, and LMH advised on formulating the research questions, designing the analysis, informing interpretation, and writing the article.

Funding NHS Digital commissioned APMS 2014 with funding from the Department of Health and Social Care in England (DHSC). The current study was part of independent research funded by the National Institute for Health Research (NIHR) Public Health Research Consortium (PHRC) Policy Research Programme ('Establishing trends and understanding context in women's mental health, 1993–2014' PHP-EHF50/27). Leonie Tanczer and Louise Howard receive salary support from the UKRI Cross-Disciplinary Network on Violence, Abuse and Mental Health (ES/S004424/1: Violence, Abuse and Mental Health). Louise Howard also receives salary support from the NIHR Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London and is an NIHR Senior Investigator. Paul Bebbington acknowledges the support of UCLH NIHR BRC. The views expressed in this article are those of the authors and not necessarily those of the NIHR or the DHSC. The funder had no role in the study design, data collection, data analysis, data interpretation, or writing of this paper. We are indebted to the thousands of people across England who generously participated in the survey.

Data availability The corresponding author (SM) had full access to all the data in the study and had final responsibility for the decision to submit for publication. The APMS dataset is lodged with the UK Data Service archive. Permission to use the dataset for this analysis was obtained from NHS Digital. Requests for further use should be made to the Data Access Request Service at NHS Digital [<https://digital.nhs.uk/services/data-access-request-service-dars>].

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Ethical approval The original survey was approved by the West London National Research Ethics Committee 14/LO/0411, RIT0985, 139324. The secondary analyses were approved by the National Centre for Social Research ethical review committee (P12395). Approval has therefore been obtained by the appropriate ethics committees and the research has been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes

were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Oram S, Boyce N, Howard LM (2019) Mental health and violence: opportunities for change. *Lancet Psychiatry* 6(7):558–559
2. Sweeney A, Clement S, Filson B, Kennedy A (2016) Trauma-informed mental healthcare in the UK: what is it and how can we further its development? *Ment Health Rev J* 21(3):174–192
3. DHSC, Agenda (2018) The Women's Mental Health Taskforce: final report. <https://www.gov.uk/government/publications/the-womens-mental-health-taskforce-report>. Accessed 26 Dec 2020
4. Wilton J, Williams A (2019) Engaging with complexity: providing effective trauma-informed care for women. CMH and MHF. https://www.centreformentalhealth.org.uk/sites/default/files/2019-04/CentreforMH_EngagingWithComplexity.pdf. Accessed 26 Dec 2020
5. AVA, Agenda (2019) Breaking down the barriers: findings of the national commission on domestic and sexual violence and multiple disadvantage. <https://avaproject.org.uk/breaking-down-the-barriers-findings-of-the-national-commission-on-domestic-and-sexual-violence-and-multiple-disadvantage/>. Accessed 26 Dec 2020
6. Scott S, Williams J, Kelly L, MacNaughton NC, Lovett J, McManus S (2013) Violence, abuse, and mental health in the English population. NatCen: London. [https://natcen.ac.uk/our-research/research/responding-effectively-to-long-term-consequences-of-violence-and-abuse-\(reva\)](https://natcen.ac.uk/our-research/research/responding-effectively-to-long-term-consequences-of-violence-and-abuse-(reva))
7. Bjørnholt M (2019) The social dynamics of revictimization and intimate partner violence: an embodied, gendered, institutional and life course perspective. *Nord J Criminol* 20(1):90–110
8. World Health Organisation (WHO) (2012) Understanding and addressing violence against women. WHO/RHR/12.36 https://apps.who.int/iris/bitstream/handle/10665/77432/WHO_RHR_12.36_eng.pdf;jsessionid=E1F9926BB8BEEB695F44CB93CD027B2?sequence=1. Accessed 26 Dec 2020
9. Criminal Prosecution Service (CPS) (2017) Controlling or coercive behaviour in an intimate or family relationship: legal guidance, domestic abuse. <https://www.cps.gov.uk/legal-guidance/controlling-or-coercive-behaviour-intimate-or-family-relationship>. Accessed 26 Dec 2020
10. Postmus JL, Hoge GL, Breckenridge J, Sharp-Jeffs N, Chung D (2020) Economic abuse as an invisible form of domestic violence: a multicountry review. *Trauma Violence Abuse* 21(2):261–283
11. UK Parliament (2021) Domestic Abuse Act 2021. <https://bills.parliament.uk/bills/2709>. Accessed 8 May 2021
12. Walby S, Towers J (2018) Untangling the concept of coercive control: theorizing domestic violent crime. *Criminol Crim Justice* 18(1):7–28
13. Yoshihama M, Horricks J, Kamano S (2009) The role of emotional abuse in intimate partner violence and health among women in Yokohama, Japan. *Am J Public Health* 99(4):647–653
14. Malicious Communications Amendment (2001) <http://www.legislation.gov.uk/ukpga/1988/27>. Accessed 26 Dec 2020
15. ONS (2020) Domestic abuse in England and Wales: year ending March 2020. <https://www.ons.gov.uk/peoplepopulationandcommunity/crimeandjustice/articles/domesticabuseprevalence>

- dtrendsenglandandwales/yearendingmarch2020. Accessed 26 Dec 2020
16. Her Majesty's Inspectorate of Constabulary and Fire and Rescue Services (2017) Living in fear—the police and CPS response to harassment and stalking. <https://www.justiceinspectors.gov.uk/hmicfrs/publications/living-in-fear-the-police-and-cps-response-to-harassment-and-stalking/>. Accessed 26 Dec 2020
 17. Leslie E, Wilson R (2020) Sheltering in place and domestic violence: Evidence from calls for service during COVID-19. *J Public Econ* 189:104241
 18. Lopez-Neira I, Patel T, Parkin S, Danezis G, Tanczer L (2019) 'Internet of things': how abuse is getting smarter. *Safe Domest Abus Q* 63:22–26
 19. Department for Digital, Culture, Media and Sport, Home Office (2019) Online harms white paper. <https://www.gov.uk/government/consultations/online-harms-white-paper>. Accessed 26 Dec 2020
 20. Barker K, Jurasz O (2017) Submission of Evidence to Scottish Government Independent Review of Hate Crime Legislation (Bracadale Review). Independent Review of Hate Crime Legislation in Scotland
 21. Livingstone S, Smith PK (2014) Annual research review: harms experienced by child users of online and mobile technologies: the nature, prevalence and management of sexual and aggressive risks in the digital age. *J Child Psychol Psychiatry* 55(6):635–654
 22. Klettke B, Hallford DJ, Clancy E, Mellor DJ, Toubmourou JW (2019) Sexting and psychological distress: the role of unwanted and coerced sexts. *Cyberpsychology Behav Soc Netw* 22(4):237–242
 23. Cox J, Koebler J, Franceschi-Bicchieri L (2018) 'Stalkerware' website let anyone intercept texts of tens of thousands of people. https://motherboard.vice.com/en_us/article/pa97g7/xnore-copy9-stalkerware-data-breach-thousands-victims. Accessed 26 Dec 2020
 24. Chatterjee R, Doerfler P, Orgad H, Havron S, Palmer J, Freed D et al (2018) The spyware used in intimate partner violence. *IEEE Symp Secur Priv*. <https://doi.org/10.1109/SP.2018.00061> (lastaccessed26/12/2020)
 25. Tanczer L, Lopez-Neira I, Parkin S, Patel T, Danezis G (2018) Gender and IoT research report: The rise of the Internet of Things and implications for technology-facilitated abuse. London: University College London. pp. 1–9. <https://www.ucl.ac.uk/steapp/sites/steapp/files/giot-report.pdf>. Accessed 26 Dec 2020
 26. UK Law Commission (2020) Harmful online communications: consultation paper 248. <https://s3-eu-west-2.amazonaws.com/lawcom-prod-storage-11jsxou24uy7q/uploads/2020/09/Online-Communications-Consultation-Paper-FINAL-with-cover.pdf>. Accessed 26 Dec 2020
 27. McManus S, Gunnell D, Cooper C, Bebbington PE, Howard LM, Brugha T, Jenkins R, Hassiotis A, Weich S, Appleby L (2019) Prevalence of non-suicidal self-harm and service contact in England, 2000–14: repeated cross-sectional surveys of the general population. *Lancet Psychiatry* 6(7):573–581
 28. McManus S, Bebbington PE, Jenkins R, Morgan Z, Brown L, Collinson D, Brugha T (2020) Data resource profile: Adult Psychiatric Morbidity Survey (APMS). *Int J Epidemiol* 49(2):361–362e
 29. McManus S, Bebbington P, Jenkins R, Brugha T (eds.) (2016) Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. Leeds: NHS Digital. <https://digital.nhs.uk/data-and-information/publications/statistical/adult-psychiatric-morbidity-survey/adult-psychiatric-morbidity-survey-of-mental-health-and-wellbeing-england-2014>. Accessed 26 Dec 2020
 30. World Health Organization (1993) The ICD-10 classification of mental and behavioural disorders: diagnostic criteria for research, vol 2. World Health Organization, Geneva
 31. Lewis G, Pelosi AJ, Araya R, Dunn G (1992) Measuring psychiatric disorder in the community; a standardised assessment for use by lay interviewers. *Psychol Med* 22:465–486
 32. Plener PL, Fegert JM (2015) Nonsuicidal self-injury: a condition for further study *Child and Adolescent Psychiatry and Mental. Health* 9:30
 33. Saunders JB, Aasland OG, Babor TF, Dela Fuente JR, Grant M (1993) Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption, part II. *Addiction* 88:791–804
 34. Straus MA (1979) Measuring intra family conflict and violence: The Conflict Tactics (CT) Scales. *J Marriage Fam* 41(1):75–88. <https://doi.org/10.2307/351733>
 35. Corp IBM, Released, (2017) IBM SPSS Statistics for Windows, Version 250. IBM Corp, Armonk
 36. StataCorp (2015) Stata Statistical Software: Release 14. Stata-Corp LP, College Station
 37. Fernet M, Lapierre A, Hébert M, Cousineau M (2019) A systematic review of literature on cyber intimate partner victimization in adolescent girls and women. *Comput Hum Behav*. <https://doi.org/10.1016/j.chb.2019.06.005>
 38. McLeod D, Flood S (2018) Coercive control: impacts on children and young people in the family environment. Dartington: Research in Practice. <https://www.researchinpractice.org.uk/children/publications/2018/december/coercive-control-impacts-on-children-and-young-people-in-the-family-environment-literature-review-2018/>. Accessed 26 Dec 2020
 39. Dymock A, van der Westhuizen C (2019) A dish served cold: targeting revenge in revenge pornography. *Leg Stud* 39(3):361–377
 40. Crofts T, Kirchengast T (2019) A ladder approach to criminalising revenge pornography. *J Crim Law* 83(1):87–103
 41. Walker K, Sleath E (2017) A systematic review of the current knowledge regarding revenge pornography and non-consensual sharing of sexually explicit media. *Aggress Violent Behav* 36:9–24
 42. Lindsay M, Booth JM, Messing JT (2016) Experiences of online harassment among emerging adults: emotional reactions and the mediating role of fear. *J Interpers Violence* 31(3):3174–3195
 43. Hobbs D (2019) Vodafone to supply mobile phones to victims of domestic abuse. Third Sector. <https://www.thirdsector.co.uk/vodafone-supply-mobile-phones-victims-domestic-abuse/digital/article/1562610>. Accessed 26 Dec 2020
 44. Herman JL (2015) Trauma and recovery: the aftermath of violence—from domestic abuse to political terror. Hachette UK
 45. Ricciardelli R, Adorjan M (2019) 'If a girl's photo gets sent around, that's a way bigger deal than if a guy's photo gets sent around': gender, sexting, and the teenage years. *J Gend Stud* 28(5):563–577
 46. De Keseredy WS, Schwartz M (1998) Measuring the extent of woman abuse in intimate heterosexual relationships: a critique of the conflict tactics scales <http://stoprelationshipabuse.org/wp-content/uploads/2013/06/A-Critique-of-the-Conflict-Tactics-Scales.pdf>. Accessed 26 Dec 2020
 47. Trevillion K, Oram S, Feder G, Howard LM (2012) Experiences of domestic violence and mental disorders: a systematic review and meta-analysis. *PLoS ONE* 7(12):e51740
 48. ONS (2015) 2011 Census analysis: what does the 2011 census tell us about people living in communal establishments?
 49. Freed D, Palmer J, Minchala D, Levy K, Ristenpart T, Dell N (2018) 'A Stalker's paradise': how intimate partner abusers

- exploit technology CHI 2018, Montréal, QC, Canada. <https://dl.acm.org/citation.cfm?id=3174241>
50. Pierce M, Hope H, Ford T, Hatch S, Hotopf M et al (2020) Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *Lancet Psychiatry* 7(10):883–892
51. Gibson J (2020) Domestic violence during COVID-19: the GP role. *Br J Gen Pract* 70(696):340
52. Pierce M, McManus S, Hope H, Hotopf M, Ford T, Hatch SL, John A, Kontopantelis E, Webb RT, Wessely R, Abel KM (2021) Mental health responses to the COVID-19 pandemic: a latent class trajectory analysis using longitudinal UK data. *Lancet Psychiatry*. [https://doi.org/10.1016/S2215-0366\(21\)00151-6](https://doi.org/10.1016/S2215-0366(21)00151-6)
53. NatCen Social Research, University of Leicester (2018) Adult Psychiatric Morbidity Survey, 2014: special licence access. UK Data Service. SN: 8203